

# *Why We Need A New Direction In Climate Change Science*



Roger A. Pielke Sr.  
Senior Research Scientist  
CIRES, University of Colorado at Boulder

Presented at the Department of Geological Sciences  
University of Texas at Austin

January 23, 2007

- The Climate System is much more than long-term weather statistics.
- Climate is not a boundary value problem but an initial value problem.
- Skillful multi-decadal climate predictions on the global, zonally-averaged and regional scales have not been achieved.
- The global-averaged surface temperature trends assessment is an inadequate climate change metric.

# *The Climate*



***Climate system:*** The system consisting of the atmosphere, hydrosphere, lithosphere, and biosphere, determining the Earth's climate as the result of mutual interactions and responses to external influences (forcing). Physical, chemical, and biological processes are involved in the interactions among the components of the climate system.

From: National Research Council, 2005: Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties, Committee on Radiative Forcing Effects on Climate, Climate Research Committee, 224 pp.

<http://www.nap.edu/catalog/11175.html>




FIGURE 1-1 The climate system, consisting of the atmosphere, oceans, land, and cryosphere. Important state variables for each sphere of the climate system are listed in the boxes. For the purposes of this report, the Sun, volcanic emissions, and human-caused emissions of greenhouse gases and changes to the land surface are considered external to the climate system.

From: National Research Council, 2005: Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties, Committee on Radiative Forcing Effects on Climate, Climate Research Committee, 224 pp.

<http://www.nap.edu/catalog/11175.html>




FIGURE 1-2 Conceptual framework of climate forcing, response, and feedbacks under present-day climate conditions. Examples of human activities, forcing agents, climate system components, and variables that can be involved in climate response are provided in the lists in each box.

From: National Research Council, 2005: Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties, Committee on Radiative Forcing Effects on Climate, Climate Research Committee, 224 pp.

<http://www.nap.edu/catalog/11175.html>

# *EXPANDING THE RADIATIVE FORCING CONCEPT (NRC 2005 Recommendations)*

- **Account for the Vertical Structure of Radiative Forcing**
- **Determine the Importance of Regional Variation in Radiative Forcing**
- **Determine the Importance of Nonradiative Forcings**
- **Provide Improved Guidance to the Policy Community**

# *Account for the Vertical Structure of Radiative Forcing*

## National Research Council Report PRIORITY RECOMMENDATIONS

- Test and improve the ability of climate models to reproduce the observed vertical structure of forcing for a variety of locations and forcing conditions.
- Undertake research to characterize the dependence of climate response on the vertical structure of radiative forcing.
- Report global mean radiative forcing at both the surface and the top of the atmosphere in climate change assessments.

# *Determine the Importance of Regional Variation in Radiative Forcing*

## National Research Council Report PRIORITY RECOMMENDATIONS:

- Use climate records to investigate relationships between regional radiative forcing (e.g., land use or aerosol changes) and climate response in the same region, other regions, and globally.
- Quantify and compare climate responses from regional radiative forcings in different climate models and on different timescales (e.g., seasonal, interannual), and report results in climate change assessments.

# *Determine the Importance of Nonradiative Forcings*

## National Research Council Report PRIORITY RECOMMENDATIONS

- Improve understanding and parameterizations of aerosol-cloud thermodynamic interactions and land-atmosphere interactions in climate models in order to quantify the impacts of these nonradiative forcings on both regional and global scales.
- Develop improved land-use and land-cover classifications at high resolution for the past and present, as well as scenarios for the future.


# *Provide Improved Guidance to the Policy Community*

## National Research Council Report PRIORITY RECOMMENDATIONS

- Encourage policy analysts and integrated assessment modelers to move beyond simple climate models based entirely on global mean TOA radiative forcing and incorporate new global and regional radiative and nonradiative forcing metrics as they become available.

# *The Narrow Focus of the IPCC Assessment*

## The global mean radiative forcing of the climate system for the year 2000, relative to 1750



Estimated radiative forcings since preindustrial times for the Earth and Troposphere system (TOA radiative forcing with adjusted stratospheric temperatures). The height of the rectangular bar denotes a central or best estimate of the forcing, while each vertical line is an estimate of the uncertainty range associated with the forcing guided by the spread in the published record and physical understanding, and with no statistical connotation. Each forcing agent is associated with a level of scientific understanding, which is based on an assessment of the nature of assumptions involved, the uncertainties prevailing about the processes that govern the forcing, and the resulting confidence in the numerical values of the estimate. On the vertical axis, the direction of expected surface temperature change due to each radiative forcing is indicated by the labels “warming” and “cooling.” From: IPCC 2001: Summary for Policymakers. A Report of the Working Group 1 of the Intergovernmental Panel on Climate Change.

<http://www.ipcc.ch/pub/spm22-01.pdf>




FIGURE 1-4 Conceptual framework for how radiative forcing fits into the climate policy framework. Blue-shaded boxes indicate quantities that have been considered as policy targets in international negotiations and other policy analyses. Radiative forcing (striped box) has not been treated as a policy target in the same explicit way that limiting emissions (e.g., Kyoto Protocol), limiting concentrations (e.g., greenhouse gas stabilization scenarios), and limiting temperature changes and impacts (e.g., environmental scenarios) have. That is, an explicit cap on anthropogenic radiative forcing levels has not been proposed analogous, for example, to the Kyoto Protocol cap on emissions. Note that land-use change has not received much attention as a forcing agent and is not included here, though this report recommends that it should be.

From: National Research Council, 2005: Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties, Committee on Radiative Forcing Effects on Climate, Climate Research Committee, 224 pp.

<http://www.nap.edu/catalog/11175.html>

# *What Fraction of Global Warming is Due to the Radiative Forcing of Increased Atmospheric Concentrations of CO<sub>2</sub>?*

## IPCC View:

- 58% of the radiative forcing of well-mixed greenhouse gases result from CO<sub>2</sub>.
- 48% of the human-caused warming climate forcing result from the radiative effect of CO<sub>2</sub>.

# *New Findings*

- i) Ozone was responsible for one-third to one-half of the observed warming trend in the Arctic during winter and spring [Drew Shindell].
- ii) The new interpretations reveal methane emissions may account for a third of the climate warming from well-mixed greenhouse gases between the 1750s and today.  
[Drew Shindell and colleagues; Keppler et al.]
- iii) For the period 2000-2004, a CERES Science Team assessment of the shortwave albedo found a decrease by 0.0015 which corresponds to an extra  $0.5 \text{ W m}^{-2}$  of radiative imbalance according to their assessment.  
[CIRES Science Team]

# *New Findings*


- iv) Model results indicate radiative forcings of  $+0.3 \text{ W m}^{-2}$  in the Northern Hemisphere associated with albedo effects of soot on snow and ice [Hansen and Nazarenko 2004]
- v) There are a variety of direct and indirect aerosol effects that cause global warming including the black carbon direct effect, the semidirect effect, and the glaciation indirect effect, with the thermodynamic effect having an unknown influence (NRC 2005).
- (these findings are summarized at <http://climatesci.atmos.colostate.edu/2006/04/27/what-fraction-of-global-warming-is-due-to-the-radiative-forcing-of-increased-atmospheric-concentrations-of-co2/>)

# *New Relative Contribution Percent of the Radiative Effect of CO<sub>2</sub>*

In Watts per meter squared


- Methane +0.8
- Short-wave albedo change +0.5
- Tropospheric ozone +0.3
- Aerosol black carbon +0.2
- Black carbon on snow and ice +0.3
- Semi-direct aerosol effect +0.1
- Glaciation effect +0.1
- Solar influences +0.25

The CO<sub>2</sub> contribution to the radiative warming decreases to 26.5% using the IPCC framework.



Multi-Decadal Land Surface  
Air Temperature Trends are  
Not A Robust Measure Of  
Global Warming and Cooling

Observed time evolution of vertical potential temperature. Note large vertical gradients near the surface [after Acevedo and Fitzjarrald, 2004].




From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. *J. Geophys. Research*, submitted.

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>

		Wind Speed (m/s)									
		10	9	8	7	6	5	4	3	2	1
Z(m)	10	0.28	0.30	0.33	0.36	0.40	0.45	0.52	0.63	0.81	1.20
	9	0.28	0.30	0.33	0.36	0.40	0.45	0.53	0.64	0.83	1.24
	8	0.28	0.30	0.33	0.36	0.40	0.46	0.53	0.65	0.85	1.28
	7	0.28	0.31	0.33	0.37	0.41	0.46	0.54	0.66	0.86	1.32
	6	0.28	0.31	0.33	0.37	0.41	0.47	0.55	0.67	0.88	1.37
	5	0.29	0.31	0.34	0.37	0.41	0.47	0.55	0.68	0.89	1.41
	4	0.29	0.31	0.34	0.37	0.42	0.48	0.56	0.69	0.91	1.46
	3	0.29	0.31	0.34	0.38	0.42	0.48	0.57	0.70	0.93	1.50
	2	0.29	0.31	0.34	0.38	0.42	0.49	0.57	0.71	0.95	1.55
	1	0.29	0.32	0.35	0.38	0.43	0.49	0.58	0.72	0.97	1.60
	0	0.29	0.32	0.35	0.38	0.43	0.50	0.59	0.73	0.98	1.66

Potential temperature increase at different levels from the experiment with  $-49 \text{ W m}^{-2}$  cooling to the experiment with  $-50 \text{ W m}^{-2}$  cooling. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>



Potential temperature increase at different levels from the experiment at  $-49 \text{ W m}^{-2}$  to the experiment with  $-50 \text{ W m}^{-2}$  cooling. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted. <http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>



**USHCN station exposure at Greensburg, Kentucky. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.**

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>



**USHCN station exposure at Leitchfield\_2\_N, Kentucky. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.**

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>




**USHCN station exposure at Leitchfield\_2\_N, Kentucky. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.**

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>



**USHCN station exposure at Hopkinsville, Kentucky. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.**

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>



**Trends of 10-year running windows for the NARR temperature anomalies. From: Pielke Sr. et al., 2006: Unresolved issues with the assessment of multi-decadal global land surface temperature trends. J. Geophys. Research, submitted.**

<http://blue.atmos.colostate.edu/publications/pdf/R-321.pdf>

**Table 5.** Linear regression trends of the monthly mean temperature anomalies (Unit: °C/10 years). The bold number indicates the trend is significant at the 5% level. The trend with superscript “n” (n = 1, 2, 3, 4, representing the column index) indicates the difference between that trend and the other trend at the n<sup>th</sup> column is significant at the 5% level (superscripts are only assigned to those trends that are significant at the 5% level).

Station	Raw Obs.	Adj. Obs.	NNR*	NARR
Trinidad	<b>0.63</b> <sup>3,4</sup>	<b>0.63</b> <sup>3,4</sup>	0.23	0.23
Cheyenne Wells	<b>0.47</b> <sup>2,4</sup>	0.33	<b>0.47</b> <sup>2,4</sup>	0.28
Las Animas	0.30	<b>0.66</b> <sup>1,3,4</sup>	<b>0.41</b> <sup>1,2,4</sup>	0.33
Eads	0.19	0.26	<b>0.49</b> <sup>1,2,4</sup>	0.28
Lamar	-0.16	<b>0.42</b> <sup>1,4</sup>	<b>0.43</b> <sup>1,4</sup>	0.29

\* All trends are calculated for the period of 1979-2003 except NNR whose trend is from 1979-1999 because our collection of the NNR data does not include any data after 1999.

From: Pielke Sr., R.A, C. Davey, J. Angel, O. Bliss, M. Cai, N. Doesken, S. Fall, K. Gallo, R. Hale, K.G. Hubbard, H. Li, X. Lin, J. Nielsen-Gammon, D. Niyogi, and S. Raman, 2006: Documentation of bias associated with surface temperature measurement sites. Bull. Amer. Meteor. Soc., submitted. <http://blue.atmos.colostate.edu/publications/pdf/R-318.pdf>

*Ocean Heat Content Changes Over Time Provides  
An Effective Metric To Diagnose The Radiative Imbalance  
of the Climate System*




Figure 1. Globally averaged annual OHCA [ $10^{22}$  J] in the upper 750 m estimated using in situ data alone from 1993 through 2005 (black line) and using in situ data excluding profiling floats (gray line). Error bars (from Figure 3) reflect the standard error estimates discussed in Section 3. Linear trends are computed from a weighted least square fit [Wunsch, 1996] and reflect the OHCA estimate made using all available profile data. Errors for inset linear trend estimates are quoted at the 95% confidence interval.

From Lyman, J.M., J. Willis, and G. Johnson, 2006: Recent cooling of the upper ocean. *Geophys Res. Letts.*, in press.





Figure 4. Globally averaged ocean temperature change [°C] from 2003 to 2005 versus depth [m]. Thin black lines represent error bounds determined by scaling the uncertainty in heat content using regression coefficients [Willis *et al.*, 2004].


From Lyman, J.M., J. Willis, and G. Johnson, 2006: Recent cooling of the upper ocean. *Geophys Res. Letts.*, in press.

# Current SST Anomalies

NOAA Current SST Anomalies (C), 1/15/2007  
(white regions indicate sea-ice)



-5.0 -4.5 -4.0 -3.5 -3.0 -2.5 -2.0 -1.5 -1.0 -0.5 0.0 0.5 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0



<http://www.osdpd.noaa.gov/PSB/EPS/SST/climo.html>

# *New or Under-Recognized Human Climate Forcings*

- Biogeochemical Effect of CO<sub>2</sub>
- Nitrogen Deposition
- Land-Use/Land-Cover Change
- Glaciation Effect of Aerosols
- Thermodynamic Effect of Aerosols
- Surface Energy Budget Effect

**TABLE 2-2 Overview of the Different Aerosol Indirect Effects Associated with Clouds**

Effect	Cloud Type	Description	Sign of TOA Radiative Forcing
First indirect aerosol effect (cloud albedo or Twomey effect)	All clouds	For the same cloud water or ice content, more but smaller cloud particles reflect more solar radiation	Negative
Second indirect aerosol effect (cloud lifetime or Albrecht effect)	All clouds	Smaller cloud particles decrease the precipitation efficiency, thereby prolonging cloud lifetime	Negative
Semidirect effect	All clouds	Absorption of solar radiation by soot leads to evaporation of cloud particles	Positive
Glaciation indirect effect	Mixed-phase clouds	An increase in ice nuclei increases the precipitation efficiency	Positive
Thermodynamic effect	Mixed-phase clouds	Smaller cloud droplets inhibit freezing, causing supercooled droplets to extend to colder temperatures	Unknown
Surface energy budget effect	All clouds	The aerosol-induced increase in cloud optical thickness decreases the amount of solar radiation reaching the surface, changing the surface energy budget	Negative

From: National Research Council, 2005: Radiative Forcing of Climate Change: Expanding the Concept and Addressing Uncertainties, Committee on Radiative Forcing Effects on Climate, Climate Research Committee, 224 pp.



<http://www.nap.edu/catalog/11175.html>

# *New Climate Change Metrics Are Needed*

- Gradient of Radiative Forcing
  - Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974. <http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>
- Change in Portioning Between Latent and Sensible Surface Heat Fluxes
  - Chase, T.N., R.A. Pielke, T.G.F. Kittel, R.R. Nemani, and S.W. Running, 2000: Simulated impacts of historical land cover changes on global climate in northern winter. *Climate Dynamics*, 16, 93-105. <http://blue.atmos.colostate.edu/publications/pdf/R-214.pdf>
- Change in Global and Regional Water Cycle
  - Marshall, C.H., R.A. Pielke Sr., and L.T. Steyaert, 2004: Has the conversion of natural wetlands to agricultural land increased the incidence and severity of damaging freezes in south Florida? *Mon. Wea. Rev.*, 132, 2243-2258. <http://blue.atmos.colostate.edu/publications/pdf/R-281.pdf>
  - Pielke, R.A. Sr., and T.N. Chase, 2003: A proposed new metric for quantifying the climatic effects of human-caused alterations to the global water cycle. Preprints, Symposium on Observing and Understanding the Variability of Water in Weather and Climate, 83rd AMS Annual Meeting, February 9-13, Long Beach, CA. <http://blue.atmos.colostate.edu/publications/pdf/PPR-249.pdf>

*Change in Portioning Between Latent  
and Sensible Surface Heat Fluxes*


# Effect of Land-Use Change on Deep Cumulonimbus Convection




From Pielke Sr., R.A., 2001: Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall. *Rev. Geophys.*, 39,151-177.

<http://blue.atmos.colostate.edu/publications/pdf/R-231.pdf>

## NATURAL VEGETATION TYPE



## CURRENT VEGETATION




Vegetation classifications for (a) natural vegetation and (b) current vegetation in regions where current and natural vegetation differ (i.e., anthropogenically disturbed regions in the current case).


From: Chase, T.N., R.A. Pielke, T.G.F. Kittel, R.R. Nemani, and S.W. Running, 2000: Simulated impacts of historical land cover changes on global climate in northern winter. *Climate Dynamics*, 16, 93-105.  
<http://blue.atmos.colostate.edu/publications/pdf/R-214.pdf>

From: Chase, T.N., R.A. Pielke, T.G.F. Kittel, R.R. Nemani, and S.W. Running, 2000: Simulated impacts of historical land cover changes on global climate in northern winter. *Climate Dynamics*, 16, 93-105. <http://blue.atmos.colostate.edu/publications/pdf/R-214.pdf>

### EAST-WEST WIND DIFFERENCE (200 mb)



### 200mb HEIGHT GRADIENT





b

Fig. 9 a Difference in 200 hPa east-west wind (current-natural), contours are  $2 \text{ m s}^{-1}$ . Shaded regions as in Fig. 3. b Comparison of north-south derivative of zonally averaged 200 hPa heights ( $d(Z_{200})/dy$ ) in Northern Hemisphere

$$Q_N + Q_H + Q_{LE} + Q_G = 0$$

$$Q_N = Q_S(1 - A) + Q_{LW}^{\downarrow} - Q_{LW}^{\uparrow}$$

From Pielke Sr., R.A., G. Marland, R.A. Betts, T.N. Chase, J.L. Eastman, J.O. Niles, D. Niyogi, and S. Running, 2002: The influence of land-use change and landscape dynamics on the climate system-relevance to climate change policy beyond the radiative effect of greenhouse gases. *Phil. Trans. A. Special Theme Issue*, 360, 1705-1719. <http://blue.atmos.colostate.edu/publications/pdf/R-258.pdf>




**The ten-year average absolute-value change in surface latent turbulent heat flux in  $W m^{-2}$  at the locations where land-use change occurred for (a) January, and (b) July.**


From Pielke Sr., R.A., G. Marland, R.A. Betts, T.N. Chase, J.L. Eastman, J.O. Niles, D. Niyogi, and S. Running, 2002: The influence of land-use change and landscape dynamics on the climate system- relevance to climate change policy beyond the radiative effect of greenhouse gases. *Phil. Trans. A. Special Theme Issue*, 360, 1705-1719.

<http://blue.atmos.colostate.edu/publications/pdf/R-258.pdf>

(a)



(b)



**The ten-year average absolute-value change in surface latent turbulent heat flux in  $\text{W m}^{-2}$  worldwide as a result of the land-use changes for (a) January, and (b) July. (Adapted from Chase et al. 2000.)**

From Pielke Sr., R.A., G. Marland, R.A. Betts, T.N. Chase, J.L. Eastman, J.O. Niles, D. Niyogi, and S. Running, 2002: The influence of land-use change and landscape dynamics on the climate system-relevance to climate change policy beyond the radiative effect of greenhouse gases. *Phil. Trans. A. Special Theme Issue*, 360, 1705-1719.

<http://blue.atmos.colostate.edu/publications/pdf/R-258.pdf>

# *Redistribution of Heat Due to the Human Disturbance of the Earth's Climate System*

Globally-Average Absolute Value of Sensible Heat Plus Latent Heat		
Only Where Land Use Occurred	July	1.08 Watts m <sup>-2</sup>
	January	0.7 Watts m <sup>-2</sup>
Teleconnections Included	July	8.90 Watts m <sup>-2</sup>
	January	9.47 Watts m <sup>-2</sup>

***Global redistribution of heat is on the same order as an El Niño.***

# *Change in the Global Water Cycle*



# *Spatial Redistribution of Heat is also Associated with a Spatial Redistribution of Water*

$$R_N = Q_G + H + L(E+T)$$
$$P = E + T + RO + I$$


*New Metric: Changes in  $\delta P$ ;  $\delta T$ ;  $\delta RO$ ;  $\delta I$*

From Pielke Sr., R.A., 2001: Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall. *Rev. Geophys.*, 39,151-177.

<http://blue.atmos.colostate.edu/publications/pdf/R-231.pdf>

# PRECIP DIFFERENCE (mm/day)


CURRENT - NATURAL



**Absolute Value of Globally-Averaged Change is 1.2 mm/day.**


# MOISTURE FLUX DIFFERENCE (mm/day)

CURRENT - NATURAL



Absolute Value of Globally-Averaged Change is 0.6 mm/day

# *Change In Regional Water Cycle*



**U.S. Geological Survey land-cover classes for pre-1900's natural conditions (left) and 1993 land-use patterns (right).**

**From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52.**

<http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>




FIG. 25. Regional average time series of accumulated convective rainfall (cm) from 1924 to 2000, with corresponding trend based on linear regression of all July-August amounts. The vertical bars overlain on the raw time series indicate the value of the standard error of the July-August regional mean.

From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52. <http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>





FIG. 26. Same as in Figure 25, except for daily (a) maximum and (b) minimum shelter-level temperature (°C)

From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52. <http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>




Associated convective rainfall (mm) from the model simulations of July-August 1973 with pre-1900s land cover (top), 1993 land use (middle), and the difference field for the two (bottom; 1993 minus pre-1900s case). From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52.

<http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>




Same as previous figure except for July and August, 1989. From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52.

<http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>






**Two-month average of the daily maximum shelter-level temperature ( $^{\circ}\text{C}$ ) from the model simulations of Jul-Aug 1989 with (top) natural land cover, (middle) current land cover. From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52.**

**<http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>**



Same as previous figure except for daily minimum temperature. From Marshall, C.H. Jr., R.A. Pielke Sr., L.T. Steyaert, and D.A. Willard, 2004: The impact of anthropogenic land-cover change on the Florida peninsula sea breezes and warm season sensible weather. *Mon. Wea. Rev.*, 132, 28-52.

<http://blue.atmos.colostate.edu/publications/pdf/R-272.pdf>



# *Gradient of Radiative Forcing*

*The Normalized Gradient of Radiative Forcing (NGoRF) is the fraction of the present Earth's heterogeneous insolation attributed to human activity on different horizontal scales*

$$NGoRF = \frac{GoRF_{anthro}}{GoRF_{total}}$$

$$GoRF_{total} = \overline{\frac{\partial R_{total}}{\partial \lambda}}$$

$$GoRF_{anthro} = \overline{\frac{\partial R_{anthro}}{\partial \lambda}}$$

From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. Geophys. Res. Letts., 33, L11813, doi:10.1029/2006GL025974.

<http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>




Figure 1. Shortwave aerosol direct radiative forcing (ADRF) for top-of atmosphere (TOA), surface, and atmosphere. From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974.

<http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>




Figure 2. Vertical profile of atmospheric heating rate ( $\text{K day}^{-1}$ ) due to shortwave ADRF. Vertical coordinate is pressure level (mb). From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974.

<http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>




Figure 3. Shortwave aerosol indirect radiative forcing (AIRF) for top-of atmosphere (TOA), surface, and atmosphere. From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974.

<http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>




Figure 4. Comparison of Mean TOA radiative forcing between infrared GRF, shortwave ADRF, and shortwave AIRF. From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974.

<http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>






Figure 5. Comparison of the meridional and the zonal component of NGoRF between infrared GRF, shortwave ADRF, and shortwave AIRF for atmosphere and surface. From: Matsui, T., and R.A. Pielke Sr., 2006: Measurement-based estimation of the spatial gradient of aerosol radiative forcing. *Geophys. Res. Letts.*, 33, L11813, doi:10.1029/2006GL025974. <http://blue.atmos.colostate.edu/publications/pdf/R-312.pdf>

# *A Focus on Vulnerability*






Schematic of the relation of water resource vulnerability to the spectrum of the environmental forcings and feedbacks (adapted from [3]). The arrows denote nonlinear interactions between and within natural and human forcings. From: Pielke, R.A. Sr., 2004: Discussion Forum: A broader perspective on climate change is needed. IGBP Newsletter, 59, 16-19.

<http://blue.atmos.colostate.edu/publications/pdf/NR-139.pdf>

## Statewide Snowpack

April 1



April 1 snowpack percent of average for the state of Colorado for years 1968 through 2006.

<ftp://ftp-fc.sc.egov.usda.gov/CO/Snow/snow/watershed/monthly/aprstatetime.gif>

# Resource Specific Impact Level with Respect to Water Resources - June 2004

## Resource Specific Impact Level Examples from Larimer County

Negligible

Minor

Moderate

Major

Exceptional



### Impacted Groups

-  Anheuser-Busch
-  Fort Collins Municipal Water
-  Grant Family Farms
-  Dryland Ranching

## ➤ *Question*

If you were given 100 million dollars to spend on environmental benefits in Colorado, where would you use that money?

1. subsidies for alternative energy
2. purchasing wilderness areas (e.g., through the Nature Conservancy)
3. building/enlarging water impoundments
4. building pipelines to transport water over large distances
5. purchasing open spaces in growing urban areas
6. funding additional mass transit


## ➤ *Where Should This Money Come From?*

1. carbon usage tax
2. mileage driven tax
3. lottery
4. tax on large private vehicles
5. state income tax increase
6. property tax increase

# *Conclusions*

The needed focus for the study of climate change and variability is on the regional and local scales. Global and zonally-averaged climate metrics would only be important to the extent that they provide useful information on these space scales.


Global warming is not equivalent to climate change. Significant, societally important climate change, due to both natural- and human- climate forcings, can occur without any global warming or cooling.



The spatial pattern of ocean heat content change is the appropriate metric to assess climate system heat changes including global warming and cooling.

Global and zonally-averaged surface temperature trend assessments, besides having major difficulties in terms of how this metric is diagnosed and analyzed, do not provide significant information on climate change and variability on the regional and local scales.

In terms of climate change and variability on the regional and local scale, the IPCC Reports, the CCSP Report on surface and tropospheric temperature trends, and the U.S. National Assessment have overstated the role of the radiative effect of the anthropogenic increase of CO<sub>2</sub> relative to the role of the diversity of other human climate forcing on global warming, and more generally, on climate variability and change.



Global and regional climate models have not demonstrated skill at predicting climate change and variability on multi-decadal time scales.

Attempts to significantly influence regional and local-scale climate based on controlling CO<sub>2</sub> emissions alone is an inadequate policy for this purpose.

A vulnerability paradigm, focused on regional and local societal and environmental resources of importance, is a more inclusive, useful, and scientifically robust framework to interact with policymakers, than is the focus on global multi-decadal climate predictions which are downscaled to the regional and local scales. The vulnerability paradigm permits the evaluation of the entire spectrum of risks associated with different social and environmental threats, including climate variability and change.

Humans are significantly altering the global climate, but in a variety of diverse ways beyond the radiative effect of carbon dioxide. The IPCC assessments have been too conservative in recognizing the importance of these human climate forcings as they alter regional and global climate. These assessments have also not communicated the inability of the models to accurately forecast the spread of possibilities of future climate. The forecasts, therefore, do not provide any skill in quantifying the impact of different mitigation strategies on the actual climate response that would occur.

# Roger A. Pielke Sr. Research Group Weblog

<http://climatesci.atmos.colostate.edu>

## Roger A. Pielke Sr. Website

<http://cires.colorado.edu/science/groups/pielke>



**PowerPoint Presentation Prepared by  
Dallas Jean Staley  
Research Assistant and Webmaster  
University of Colorado  
Boulder, Colorado 80309**